

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/115374/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Thiry, Valentine, Clauss, Marcus, Stark, Danica, Beudels-Jamar, Roseline C., Vercauteren Drubbel, Régine, Nathan, Senthilvel K.S.S., Goossens, Benoit
ORCID: <https://orcid.org/0000-0003-2360-4643> and Vercauteren, Martine
2018. Faecal particle size in free-ranging proboscis monkeys, *Nasalis larvatus*: variation between seasons. *Folia Primatologica* 89 (5) , pp. 327-334.
10.1159/000490794 file

Publishers page: <http://dx.doi.org/10.1159/000490794>
<<http://dx.doi.org/10.1159/000490794>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Title:

Faecal particle size in free-ranging proboscis monkeys, *Nasalis larvatus*: Variation between seasons

Authors:

Valentine Thiry^{ab}, Marcus Clauss^c, Danica J. Stark^{de}, Roseline C. Beudels-Jamar^b,
Régine Vercauteren Drubbel^a, Senthilvel K. S. S. Nathan^f, Benoît Goossens^{defg},
Martine Vercauteren^a

^a Anthropology and Human Genetics Unit, Université libre de Bruxelles, Brussels, Belgium;

^b Conservation Biology Unit, Education and Nature, Royal Belgian Institute of Natural Sciences, Brussels, Belgium;

^c Clinic for Zoo Animals, Exotic Pets and Wildlife, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland;

^d Organisms and Environment Division, Cardiff School of Biosciences, Cardiff University, Sir Martin Evans Building, Museum Avenue, Cardiff CF10 3AX, United Kingdom;

^e Danau Girang Field Centre c/o Sabah Wildlife Department, Wisma Muis, Kota Kinabalu, Sabah 88100, Malaysia;

^f Sabah Wildlife Department, Kota Kinabalu, Sabah, Malaysia;

^g Sustainable Places Research Institute, Cardiff University, 33 Park Place, Cardiff CF10 3BA, UK

Short title: Faecal particle size in proboscis monkeys

Corresponding author: Valentine Thiry

Postal address : 64, Avenue des Grenadiers, 1050 Brussels, Belgium.

Phone : 0032473563831

e-mail : valentinethiry89@gmail.com

Abstract

Reducing the size of food particles is crucial for herbivores. Seasonal dietary changes are known to influence animals' chewing efficiency. Proboscis monkeys (*Nasalis larvatus*) are foregut fermenters, with a high chewing efficiency allowing them to achieve very fine faecal particles. In this study, we investigated how proboscis monkeys' chewing efficiency varies among wet and dry seasons, hypothesising differences possibly related to diet change. Faecal particle size analysis is an established approach to estimate chewing efficiency in mammalian herbivores. We analysed 113 proboscis monkey faecal samples collected in the Lower Kinabatangan Wildlife Sanctuary, between 2015 and 2017. By following standard sieve analysis protocols, we measured a mean particle size $MPS_{(0.025-8)}$ of 0.45 ± 0.14 mm, and confirmed a previous result that proboscis monkeys have a very low faecal MPS. This study highlights a seasonal influence on proboscis monkeys' chewing efficiency, with smaller MPS (better chewing efficiency) during the wet season. During that time of the year, individuals may potentially change their diet, as all faecal samples contained intact seeds. Whether the seasonal MPS difference in proboscis monkeys is smaller than in other colobines due to their 'rumination' strategy remains to be investigated.

Keywords

Nasalis larvatus, chewing efficiency, nutritional ecology, forestomach, diet change, foregut fermenter

Number of words in the manuscript: 3365

Introduction

Reducing the size of food particles is crucial in herbivores that rely on gut microbiota to digest plant components. In herbivores, fibre digestion relies on digesta retention and particle size. To accomplish the same digestibility, large digesta particles will need longer retention times than smaller ones [Bjorndal et al., 1990]. Measuring faecal particle size by wet sieving analysis, is an established non-invasive approach to determine the chewing efficiency of mammals [Fritz et al., 2009]. Several studies focused on faecal particle size in mammals [Fritz et al., 2009; Clauss et al., 2015], or more specifically in ruminants [Renecker and Hudson, 1990; Clauss et al., 2002] and primates [Dunbar and Bose, 1991; Matsuda et al., 2014; Venkataraman et al., 2014; Weary et al., 2017]. Across mammals, the size of faecal particles usually increases with animal body mass [Fritz et al., 2009]. However, among primates, the proboscis monkey (*Nasalis larvatus*) displays a particularly small mean particle size (MPS) for its average body mass (15 kg) [Matsuda et al., 2014]. Proboscis monkeys are foregut fermenters [Matsuda et al., 2014]. Like other colobine primates, they have a sacculated forestomach where the food is fermented [Bauchop, and Martucci, 1968; Milton, 1993]. Regurgitation and remastication (i.e. rumination) has been observed in wild proboscis monkeys [Matsuda et al., 2011a]. Whether this facultative rumination strategy explains how proboscis monkeys achieve particularly fine faecal particles remains unclear [Matsuda et al., 2014].

Seasonal dietary change and dental wear are known to influence animals' chewing efficiency [Venkataraman et al., 2014]. Within a species, faecal particle size can vary in relation to diet [Renecker and Hudson, 1990] or seasons [Nygren and Hofmann, 1990]. For instance, in gelada baboons (*Theropithecus gelada*), chewing efficiency decreases during the dry season when individuals feed on tougher non-preferred food items, with a more distinct effect in older individuals [Venkataraman et al., 2014]. The opposite is observed in

frugivorous chimpanzees (*Pan troglodytes schweinfurthii*): MPS is higher when chimpanzees feed on drupe fruits (preferred foods) than on figs (non-preferred foods) [Weary et al., 2017]. The authors suggested that chewing efficiency might be less critical in frugivores than in typical folivores, because they did not observe an effect of age on MPS.

With their natural diet, proboscis monkeys are excellent candidates to investigate how MPS might change throughout the year. While proboscis monkeys were first considered essentially folivores, it is now recognised that they preferentially feed on unripe fruits/seeds when they are available [Matsuda et al., 2009]. The present study investigates how proboscis monkeys' chewing efficiency varies among wet and dry seasons. We hypothesised that proboscis monkeys will achieve a higher chewing efficiency (MPS will decrease) during the season when individuals are able to consume their preferred food.

Moreover, we investigated some methodological aspects of sieve analysis. Extending the sieve column (adding larger top or smaller bottom sieves) is known to influence MPS measurements [Fritz et al., 2012], as well as including the weight of unchewed items (i.e. large seeds) and maximum particle length (MPL) in the MPS calculation [Weary et al., 2017]. Therefore, we combined various sets of sieves, with or without the MPL, and assessed the impact on MPS measurements.

Material and Methods

Study site

Our study took place in the Lower Kinabatangan Floodplain (5°20'-05°45' N, 117°40'-118°30' E), in Eastern Sabah (Malaysian Borneo), between 2015 and 2017. Daily temperatures and rainfall were measured at the research station. Below we will refer to dry season (May-June-July) where the mean monthly rainfall is 120 (\pm SD=100) mm and to the

wet season (November-February) where it reaches 243 (\pm SD=104) mm. Mean minimum and maximum temperatures reached 24.4 (\pm SD=0.6) and 30 (\pm SD=1.7) °C, respectively.

Faecal sampling

In riverine forests, proboscis monkeys are known to take refuge along riverbanks to spend the night [Matsuda et al., 2011b]. During this study, we conducted boat-based surveys along the Kinabatangan River, in the late afternoon, to find proboscis monkey groups settled at their sleeping sites. To avoid sampling the same group multiple times, we searched for proboscis monkey groups in different parts (North and South riverbanks) along a pre-established 21 km transect in a month. In the morning, we travelled back to the group's location of the previous evening. Once the group left the riverside to forage further inland, we moved to the riverbank to search for fresh faecal samples that had fallen under sleeping trees. We collected large samples, presumed to belong to adult individuals (undistinguished sex). Between May and July 2015, January and February 2016, and November 2016 and February 2017, two faecal samples were collected per group and placed in separate tubes, to perform two different analyses: manual and wet sieving analyses. 137 samples (15 ± 5 samples/month) were analysed by the manual method and 113 faecal samples (13 ± 6 samples/month) by wet sieving method.

Manual analyses

Faecal samples were cleaned with water in a 0.4 mm mesh strainer to discard faecal matter. The remaining digested items were searched for intact seeds. Percentages of samples containing seeds were used to assess seasonal changes.

Wet sieving analyses

Faecal samples were stored in a tube with 70% ethanol [Matsuda et al., 2014]. They were analysed using the standard wet sieving method [Fritz et al., 2012]. Before sieving, each sample was suspended in a beaker filled with water that was stirred continuously for 12 hours. The sample was then poured over a series of 10 sieves with mesh size of 8, 4, 2, 1, 0.5, 0.25, 0.125, 0.063, 0.04 and 0.025 mm (Retsch AS 200 digit, Haan, Germany). We conducted the sieving analysis for 10 minutes with an amplitude of 2 mm and a water flow of approximately 2 l/min. If a particle was retained on the largest sieve, its size was recorded as the maximum particle size. Particles retained on each sieve were transferred onto pre-weighed petri dishes and dried at 103°C overnight. After cooling in a desiccator, petri dishes were weighed with an analysis balance with measuring accuracy of 1 mg (Kern AEJ 220-4M, Kern, Balingen, Germany). When large seeds (≥ 2 mm) were retained intact in sieves, they were removed, weighed and subtracted from the respective sieve weight. However, the smaller (< 2 mm) and numerous seeds, such as *Ficus* and *Nauclea* seeds, were logistically impossible to remove from the analysis [Weary et al., 2017].

Among various indices, the discrete mean has been proposed as a standard to describe the MPS value obtained from sieving analyses [Fritz et al., 2012]. To compare our results with a previous study conducted on proboscis monkey's faecal particle sizes [Matsuda et al., 2014], we excluded the two smallest sieves (mesh sizes: 0.040 and 0.025 mm) from the MPS calculation, as they were not used by Matsuda et al. [2014]. Although the latter study used a larger top sieve (16 mm) than we did, no particles were ever retained on it.

Statistical analyses

We carried out t-tests on log-transformed data to compare $MPS_{(0.025-8mm)}$ values between dry and wet seasons. We compared MPS values calculated for 28 faecal samples, with or without

considering the length of the maximum particles and using series of 10 (0.025-8 mm) or 8 (0.063-8 mm) sieves. To assess the difference between those four MPS measurements, we performed a related-samples Wilcoxon signed rank test between all pairs of MPS values, using the Bonferroni adjustment (for multiple comparisons). R 3.4.0 [R Development Core Team, 2016] was used for all statistical analyses, with statistical significance of $p < 0.05$.

Results

Manual analyses

By cleaning fresh faeces (N=137), we observed that the percentage of faecal samples containing intact seeds changed throughout the year (Fig. 1), with a mean of 100 ± 0 % during the wet season (November-February) and of 38 ± 10 % during the dry season (May-July).




Figure 1 Percentage of faecal samples of proboscis monkeys (*Nasalis larvatus*) collected during different months that contain intact seeds

Wet sieving analyses

Fig. 2 illustrates the typical way faecal particles are distributed after wet sieving analysis.




Figure 2. Distribution of proboscis monkey (*Nasalis larvatus*) faecal particles after wet sieving analysis. Each petri dish represents one of the 10 cascade sieves, ordered from 8 mm until 0.025 mm mesh size. The sample PMF217-1 collected on the 28th of January 2017 was used for this example.

By using a cascade of 10 sieves (mesh sizes: 8 to 0.025 mm), the MPS reaches an average of 0.45 ± 0.14 mm and increases over the course of the observation period from November to July (Fig. 3). By using a series of eight sieves (mesh sizes: 8 to 0.063 mm) like in Matsuda et al. (2014), we observe that $MPS_{(0.063-8mm)}$ is significantly larger than $MPS_{(0.025-8mm)}$, reaching 0.55 ± 0.14 mm ($V=6441$, $N=113$, $p<0.001$).




Figure 3. Variation in proboscis monkey (*Nasalis larvatus*) MPS over the course of the observation period (November-July)

When comparing $MPS_{(0.025-8mm)}$ of proboscis monkeys across seasons, we observe that MPS is significantly smaller during the wet season than the dry season ($MPS_{wet} = 0.38 \pm 0.11$ mm, $MPS_{dry} = 0.52 \pm 0.13$ mm; t-test $t = -6.2812$, $p < 0.001$) (Fig. 4).




Figure 4. MPS variation of proboscis monkeys (*Nasalis larvatus*) across seasons

Table 1 summarises four MPS measurements. Using a series of 10 sieves (with both smallest mesh sizes 0.040 and 0.025 mm) resulted in finer MPS than calculation with the 8-sieve cascade. MPS measurements were significantly larger when including the maximum particle length (MPL) for the largest sieve in the calculation.

Table 1. Mean particle size \pm SD of 28 faecal samples of proboscis monkeys (*Nasalis larvatus*) measured using two series of sieves (indicated by the sieve size of the smallest and largest sieves) and with or without taking the maximum particle length (MPL, when particles were retained on the largest sieve) into account in the MPS calculation

Method	MPS \pm SD (mm)
0.025-8 MPL	0.48 ± 0.11
0.025-8	0.47 ± 0.11
0.063-8 MPL	0.59 ± 0.11
0.063-8	0.57 ± 0.10

Discussion

This study focused on proboscis monkeys' faecal samples collected between 2015 and 2017, during wet and dry seasons. We confirm the very small discrete mean faecal particle size in proboscis monkey ($\text{MPS}_{(0.025-8\text{mm})} = 0.45 \pm 0.14$ mm), for its average body mass. We measured a $\text{MPS}_{(0.063-8\text{mm})}$ of 0.55 ± 0.14 mm, similar to the results of a previous study ($\text{MPS}_{(0.063-16\text{mm})} = 0.53 \pm 0.09$ mm) obtained by analysing 10 samples collected in June-July 2010 [Matsuda et al., 2014]. The fine MPS indicates the generally high chewing efficiency of proboscis monkeys. For example, in comparison, frugivorous primates in Borneo, such as macaques (*Macaca fascicularis* and *M. nemestrina*) and orangutans (*Pongo pygmaeus*), have larger $\text{MPS}_{(0.063-16\text{mm})}$ respectively ranging from 1.07 ± 0.47 to 2.30 ± 0.78 mm [Matsuda et al., 2014].

We observed that MPS is even smaller during the wet season than the dry season. The same pattern had been observed in folivorous gelada baboons [Venkataraman et al., 2014]. The latter showed a lower MPS in wet season when consuming less tough food items. In our study, the MPS difference might also be linked to a change in diet. Smaller MPS values during the wet season correlate with high percentage of intact seeds in faeces, suggesting

individuals might consume more fruits and their seeds. However, this assumption must be considered carefully, as the absence of seeds in faeces does not always imply that individuals did not eat fruits (i.e. seeds could be totally digested, chewed or discharged). Mismatches have been observed between proboscis monkey fruit feeding activity and seeds detected in faeces [Matsuda et al., 2013]. Primates, including proboscis monkeys, usually avoid feeding on tough leaves or leaf parts [Hill and Lucas, 1996; Teaforde et al., 2006; Dunham and Lambert, 2016; Matsuda et al., 2017]. The same pattern is observed in chimpanzees where fallback foods are significantly tougher than preferred items (fruits) [Vogel et al., 2008]. However, in Bornean orangutans (*P. p. wurmbii*), mechanical properties of leaves and fruits did not vary significantly [Vogel et al., 2008]. In the present study, we did not analyse the toughness of food items fed on by proboscis monkeys. However, in comparison to leaves, fruits and seeds are generally considered as high quality food [Milton, 1993; Hanya and Bernard, 2015]. Containing less fibre, fruits generally are more digestible than leaves [Milton, 1993]; consumption of unripe fruits may lead to smaller MPS. Further work should investigate nutritional and mechanical properties of unripe fruits and their seeds consumed by the proboscis monkey to better understand the feeding selection in this endangered primate.

As in gelada baboons, we suggest here that fallback food consumption during some parts of the year leads to a reduction of chewing efficiency which might potentially negatively impact the animals' fitness [Venkataraman et al., 2014]. There is preliminary evidence that 'rumination' activity in proboscis monkeys is higher during times of increased leaf consumption [Matsuda et al., 2014], which could potentially attenuate the change in MPS associated with leaves. If this was a general pattern, then the MPS difference between the seasons obtained in the present study should be of a lower magnitude than in other arboreal primates that show a seasonal foraging pattern but do not 'ruminate'. Compared to geladas with a seasonal MPS difference of 0.3-0.4 mm in prime adults, the proboscis monkeys of the

present study did show a lower difference (0.14 mm, Fig. 4). However, due to diet differences between hindgut fermenter geladas and foregut fermenter proboscis monkeys, this comparison should be treated cautiously. We suggest further studies should determine whether proboscis monkey individuals achieve finer MPS when they ‘ruminate’ as opposed to times when they do not. Such data could help unravel the relevance of facultative rumination as a response to diet constraints. Finally, future research should also investigate if seasonal changes in MPS are also measured in other colobine primates, as data is missing so far.

Acknowledgements

We thank the Sabah Wildlife Department and Sabah Biodiversity Centre for allowing us to carry out our research in the Lower Kinabatangan Wildlife Sanctuary. We thank our Belgian financial funders: the FNRS (Fonds de la Recherche Scientifique), FNRS Gustave Boël-Sofina fellowship 2016 and the Fonds Léopold III – pour l’Exploration et la Conservation de la Nature asbl. Thanks to all students and research assistants at Danau Girang Field Centre who helped us in the field.

References

- Bauchop T, Martucci, RW (1968). Ruminant-like digestion of the langur monkey. *Science, New Series* 161: 698–700.
- Bjorndal KA, Bolten AB, Moore JE (1990). Digestive fermentation in herbivores: Effect of food particle size. *Physiological Zoology* 63: 710–721.
- Clauss M, Lechner-Doll M, Streich WJ (2002). Faecal particle size distribution in captive wild ruminants: An approach to the browser/grazer dichotomy from the other end. *Oecologia* 131: 343–349.
- Clauss M, Steuer P, Erlinghagen-Lückerath K, Kaandorp J, Fritz J, Südekum K-H, Hummel J (2015). Faecal particle size: Digestive physiology meets herbivore diversity. *Comparative Biochemistry and Physiology A* 179: 182–191.
- Dunbar RIM, Bose U (1991). Adaptation to grass-eating in gelada baboons. *Primates* 32: 1–7.
- Dunham NT, Lambert AL (2016). The role of leaf toughness on foraging efficiency in Angola black and white colobus monkeys (*Colobus angolensis palliatus*). *American Journal of Physical Anthropology* 161: 343–354.
- Fritz J, Hummel J, Kienzle E, Arnold C, Nunn C, Clauss M (2009). Comparative chewing efficiency in mammalian herbivores. *Oikos* 118: 1623–1632.
- Fritz J, Streich WJ, Schwarm A, Clauss M (2012). Condensing results of wet sieving analyses

into a single data: A comparison of methods for particle size description. *Journal of Animal Physiology and Animal Nutrition* 96: 783–797.

Hanya G, Bernard H (2015). Different roles of seeds and young leaves in the diet of red leaf monkeys (*Presbytis rubicunda*): Comparisons of availability, nutritional properties, and associated feeding behavior. *International Journal of Primatology* 36: 177–193.

Hill DA, Lucas PW (1996). Toughness and fiber content of major leaf foods of Japanese macaques (*Macaca fuscata yakui*) in Yakushima. *American Journal of Primatology* 38: 221–231.

Matsuda I, Clauss M, Tuuga A, Sugau J, Hanya G, Yumoto T, Bernard H, Hummel J (2017). Factors affecting leaf selection by foregut-fermenting proboscis monkeys: New insight from in vitro digestibility and toughness of leaves. *Scientific Reports* DOI:10.1038/srep42774.

Matsuda I, Higashi S, Otani Y, Tuuga A, Bernard H, Corlett RT (2013). A short note on seed dispersal by colobines: The case of the proboscis monkey. *Integrative Zoology* 8:395–399.

Matsuda I, Murai T, Clauss M, Yamada T, Tuuga A, Bernard H, Higashi S (2011a). Regurgitation and remastication in the foregut-fermenting proboscis monkey (*Nasalis larvatus*). *Biology letters* 7: 786–789.

Matsuda I, Tuuga A, Bernard H (2011b). Riverine refuging by proboscis monkeys (*Nasalis larvatus*) and sympatric primates: Implications for adaptive benefits of the riverine habitat. *Mammalian Biology* 76: 165–171.

307

308 Matsuda I, Tuuga A, Hashimoto C, Bernard H, Yamagiwa J, Fritz J, Tsubokawa K, Yayota
309 M, Murai T, Iwata Y, Clauss M (2014). Faecal particle size in free-ranging primates supports
310 a “rumination” strategy in the proboscis monkey (*Nasalis larvatus*). *Oecologia* 174: 1127–
311 1137.

312

313 Matsuda I, Tuuga A, Higashi S (2009). The feeding ecology and activity budget of proboscis
314 monkeys. *American Journal of Primatology* 71: 478–492.

315

316 Milton K (1993). Diet and primate evolution. *Scientific American* 269: 86–93.

317

318 Nygren K, Hofmann, RR (1990). Seasonal variations of food particle size in Moose. *Alces*
319 26: 44–50.

320

321 R Development Core Team (2016). R: A language and environment for statistical computing.
322 R foundation for statistical computing. R 3.4.0 GUI 1.70. Vienna, Austria. [http://www.R-](http://www.R-project.org)
323 [project.org](http://www.R-project.org)

324

325 Renecker LA, Hudson RJ (1990). Digestive kinetics of moose (*Alces alces*), wapiti (*Cervus*
326 *elaphus*) and cattle. *Animal Production* 50: 51–61.

327

328 Teaforde MF, Lucas PW, Ungar PS, Glander KE (2006). Mechanical defenses in leaves eaten
329 by Costa Rican howling monkeys (*Alouatta palliata*). *American Journal of Physical*
330 *Anthropology* 129: 99–104.

331

332 Venkataraman VV, Glowacka H, Fritz J, Clauss M, Seyoum C, Nguyen N, Fashing PJ (2014).
333 Effects of dietary fracture toughness and dental wear on chewing efficiency in geladas
334 (*Theropithecus gelada*). *American Journal of Physical Anthropology* 155: 17–32.
335
336 Vogel ER, van Woerden JT, Lucas PW, Utami Atmoko SS, van Schaik CP, Dominy NJ
337 (2008). Functional ecology and evolution of hominoid molar enamel thickness: *Pan*
338 *troglodytes schweinfurthii* and *Pongo pygmaeus wurmbii*. *Journal of Human Evolution*
339 55: 60–74.
340
341 Weary TE, Wrangham RW, Clauss M (2017). Applying wet sieving fecal particle size
342 measurement to frugivores: A case study of the eastern chimpanzee (*Pan troglodytes*
343 *schweinfurthii*). *American Journal of Physical Anthropology* 163: 510–518.